



1/9



**FIG. I**


2/9



**FIG.2**


3/9

**EFFECT OF 2'-O-METHYL GAPPED OLIGOS**




**FIG.3**

4/9



**FIG.4**


5/9

**FIG.5**

6/9


FIG. 6

PLASMA CONCENTRATION OF ISIS 3082, AND 2' O-PROPYL ANALOGS  
(ISIS 9045 AND 9046) IN BALB/c MICE AFTER INTRAVENOUS INJECTION



7/9


**ISIS 3082 TISSUE DISTRIBUTION IN BALB/c MICE  
 AS PERCENT OF DOSE AT 1, 3, 8 AND 24 HOURS AFTER  
 INTRAVENOUS INJECTION**



**FIG. 7**

8/9

**ISIS 9045 (2' O-PROPYL PHOSPHOROTHIATE ANALOG OF ISIS 3082)  
 TISSUE DISTRIBUTION IN BALB/c MICE  
 AS PERCENT OF DOSE AT 1, 3, 8 AND 24 HOURS AFTER  
 INTRAVENOUS INJECTION**



**FIG. 8**



9/9

TISSUE DISTRIBUTION OF ISIS 9046 (2' O-PROPYL GAPMER) IN BALB/c MICE  
 AT 1, 3, 8 AND 24 HOURS AFTER INTRAVENOUS INJECTION




FIG.9